ERRATA SHEET: A Basic Package of Health Services for Afghanistan, 2005/1384

Please note the following changes to ensure consistency among BPHS tables and with existing national policies. Additions are shown in **bold underline** and deletions in strikethrough.

Table 8: Essential Drugs for BPHS by Type of Facility Pages 26-33

Table 8: Essential Drugs for BPHS by Type of Facility		Pages 26-33				
Drug	Dosage Form and Strength	Health Post	внс	СНС	DH	
2.1 Non-Opiod Analge						
Acetaminophen (Paracetamol)	tablet <u>500mg</u>	√	√	$\sqrt{}$	$\sqrt{}$	
3. Anticonvulsants/ Au	nti-Epileptics					
Magnesium sulphate	Injection 500 mg/ ml in 20-ml ampoule		√	$\sqrt{}$	$\sqrt{}$	
6.1 Anthelminthics						
Mebendazole	Chewable tablet 100 mg					
6.4.2 Antimalarial and	l Antileishmanial					
Chloroquine	Tablet 150 mg (as phosphate or sulfate) 5ml		$\sqrt{}$	$\sqrt{}$	$\sqrt{}$	
Pyrimethamine +						
Suphadoxine	Tablet 25mg + 500mg	$\frac{\sqrt{1}}{\sqrt{1}}$			$\sqrt{}$	
(fansidar)						
Quinine	Tablet 300mg (as bisulphate or sulphate)		$\sqrt{2}$	\checkmark	$\sqrt{}$	
Artesunate ³	Tablet 50mg		$\sqrt{4}$			
Artesunate ³	Tablet 100mg		$\sqrt{4}$			
Artemether For pre-referral						
treatment of suspected	Injection 20 & 80 mg in oil for					
and confirmed severe	intramuscular injection		_	_		
or <i>complicated</i> malaria.						
Sodium						
Stibogluconate	Injection 100mg/ml				$\sqrt{}$	
Melamine	Injection 85mg/ml				<u>√</u>	
Antimonite		<u> </u>	.1 .	1 0	C 1	

¹Presumptive treatment for unconfirmed malaria is chloroquine *and* sulfadoxine / pyrimethamine, before referral to CHC for confirmatory diagnosis.

⁴ In BHCs where diagnostic services are available.

14.2 Contraceptives								
Medroxy Progestrone	Depot injection 150 mg/ml in 1-ml		2/	2/	2/			
(DMPA)	vial		<u>_v</u>	V	V			
IUD								
21. Vitamins and Minerals								
Retinol	Sugarcoated tablet 100,000 IU (as	ما	ما	2/	2/			
	palmitate) (5.5 mg) (55 mg)	V	V	V	V			
Multi-micronutrients								

Note: These errata are already corrected in the Dari and Pushtu versions of BPHS 2005/1384 so no errata sheet is required for those versions.

² Quinine – 2nd line treatment and treatment for severe / complicated malaria require *laboratory confirmation*³ Artemesinin combination therapy: Artesunate + SP (fansidar) as first line treatment for *laboratory confirmed* falciparum malaria..

Antimalarial and Antileishmanial Treatment:

Table 14: Health Post Summary of BPHS Services: Page 40, add <u>fansidar</u> to Anti-malarial treatment essential drugs

Table 15: Basic Health Center Summary of BPHS Services: Page 44, remove quinine from Anti-malarial treatment essential drugs

Table 16: Comprehensive Health Center Summary of BPHS Services: Page 49, add artesunate to Anti-malarial treatment essential drugs

Table 17: District Hospital Summary of BPHS Services: Page 56, add <u>artesunate</u> to Antimalarial treatment essential drugs;

Page 57, add "<u>diagnosis and treatment of leishmaniasis</u>" to services provided and <u>sodium stibogluconate</u> and <u>meglumine antimoniate</u> to Anti-leishmanial treatment essential drugs

Note: These errata are already corrected in the Dari and Pushtu versions of BPHS 2005/1384 so no errata sheet is required for those versions.